

The geometry of the space of vortex-antivortex pairs

Martin Speight (Leeds)
joint with
Nuno Romão (Augsburg)
[arXiv:1807.00712](https://arxiv.org/abs/1807.00712)

Math Phys seminar, KIAS Seoul, 20/9/18

- Vortices: simplest topological solitons in gauge theory (2D, $U(1)$, \mathbb{C})
- Nice generalization: Higgs field takes values in a kähler mfd X with hamiltonian action of gauge group G
- G action on X can have more than one fixed point: more than one species of vortex
- Different species can coexist in stable equilibrium — but can't coincide
- **Noncompact** vortex moduli spaces (even on compact domains)
- Completeness? Finite volume? Curvature properties?
- Simplest version already interesting: $X = S^2$, $G = U(1)$

- Fix $\mathbf{e} \in S^2$ (e.g. $\mathbf{e} = (0, 0, 1)$)
 $G = U(1)$ acts on S^2 by rotations about \mathbf{e}
- $P \rightarrow \Sigma$ principal G bundle, degree $n \geq 0$, connexion A
- \mathbf{n} section of $P \times_G S^2$
- Canonical sections $\mathbf{n}_\infty(x) = \mathbf{e}$, $\mathbf{n}_0(x) = -\mathbf{e}$
- **Two** integer topological invariants of a section \mathbf{n} :

$$n_+ = \#(\mathbf{n}(\Sigma), \mathbf{n}_\infty(\Sigma)), \quad n_- = \#(\mathbf{n}(\Sigma), \mathbf{n}_0(\Sigma))$$

Constraint: $n = n_+ - n_-$ (so we're assuming $n_+ \geq n_-$)

- Energy

$$E = \frac{1}{2} \int_{\Sigma} (|d_A \mathbf{n}|^2 + |F_A|^2 + (\mathbf{e} \cdot \mathbf{n})^2)$$


where, in a local trivialization

$$d_A \mathbf{n} = d\mathbf{n} - A\mathbf{e} \times \mathbf{n}, \quad F_A = dA$$

Aside: $\mu(\mathbf{n}) = -\mathbf{e} \cdot \mathbf{n}$ is moment map for gauge action


(Anti)vortices

"north" vortex




$$n_+ = 1, n_- = 0$$

"north" antivortex




$$n_+ = -1, n_- = 0$$

"south" vortex



$$n_+ = 0, n_- = -1$$

"south" antivortex



$$n_+ = 0, n_- = 1$$

“Bogomol’nyi” bound (Schroers)

- Given (\mathbf{n}, A) define a two-form on Σ

$$\Omega(X, Y) = (\mathbf{n} \times d_A \mathbf{n}(X)) \cdot d_A \mathbf{n}(Y)$$

- Let $e_1, e_2 = Je_1$ be a local orthonormal frame on Σ . Then

$$\begin{aligned}\mathcal{E} &= \frac{1}{2}(|d_A \mathbf{n}(e_1)|^2 + |d_A \mathbf{n}(e_2)|^2) + \frac{1}{2}|F_A|^2 + \frac{1}{2}(\mathbf{e} \cdot \mathbf{n})^2 \\ &= \frac{1}{2}|d_A \mathbf{n}(e_1) + \mathbf{n} \times d_A \mathbf{n}(e_2)|^2 + \frac{1}{2}|F_A - * \mathbf{e} \cdot \mathbf{n}|^2 \\ &\quad + *(\Omega + \mathbf{e} \cdot \mathbf{n} F_A)\end{aligned}$$

$$\implies E \geq \int_{\Sigma} (\Omega + \mathbf{e} \cdot \mathbf{n} F_A)$$

- Claim: last integral is a homotopy invariant of (\mathbf{n}, A)

“Bogomol’nyi” bound

- Suffices to show this in case $D = \mathbf{n}^{-1}(\{\mathbf{e}, -\mathbf{e}\}) \subset \Sigma$ finite
- On $\Sigma \setminus D$ have global one-form $\xi = \mathbf{e} \cdot \mathbf{n}(A - \mathbf{n}^* d\varphi)$ s.t.

$$\Omega + \mathbf{e} \cdot \mathbf{n} F_A = d\xi$$

- Hence

$$\begin{aligned}\int_{\Sigma} (\Omega + \mathbf{e} \cdot \mathbf{n} F_A) &= \int_{\Sigma \setminus D} (\Omega + \mathbf{e} \cdot \mathbf{n} F_A) \\ &= \lim_{\varepsilon \rightarrow 0} \sum_{p \in D} - \oint_{C_\varepsilon(p)} \xi \\ &= 2\pi(n_+ + n_-)\end{aligned}$$

“Bogomol’nyi” bound

- Hence $E \geq 2\pi(n_+ + n_-)$ with equality iff

$$\bar{\partial}_A \mathbf{n} = 0 \quad (V1)$$

$$*F_A = \mathbf{e} \cdot \mathbf{n} \quad (V2)$$

- Note solutions of $(V1)$ certainly have D finite (and $n_{\pm} \geq 0$)

- If Σ compact, there's a “Bradlow” obstruction

$$2\pi(n_+ - n_-) = \int_{\Sigma} F_A = \int_{\Sigma} \mathbf{e} \cdot \mathbf{n} \leq \text{Vol}(\Sigma)$$

- **Theorem:** Let $n_+ \geq n_- \geq 0$ and $2\pi(n_+ - n_-) < \text{Vol}(\Sigma)$. For each pair of disjoint effective divisors D_+, D_- in Σ of degrees n_+, n_- there exists a unique gauge equivalence class of solutions of (V1), (V2) with $\mathbf{n}^{-1}(\pm \mathbf{e}) = D_{\pm}$.
- Moduli space of vortices: $M_{n_+, n_-} \equiv M_{n_+} \times M_{n_-} \setminus \Delta_{n_+, n_-}$
- If $n_- > 0$, M_{n_+, n_-} is noncompact (in an interesting way)

The “Taubes” equation

$$u = \frac{n_1 + in_2}{1 + n_3}, \quad h = \log |u|^2, \quad g_\Sigma = \Omega(z) dz d\bar{z}$$

- h finite except at \pm vortices, $h = \mp\infty$.
- $(V1) \Rightarrow A_{\bar{z}} = -i \frac{\partial_{\bar{z}} u}{u}$, eliminate A from $(V2)$

$$\nabla^2 h - 2\Omega \tanh \frac{h}{2} = 0$$

away from vortex positions

- (+) vortices at z_r^+ , $r = 1, \dots, n_+$, (-) vortices at z_r^- ,
 $r = 1, \dots, n_-$

$$\nabla^2 h - 2\Omega \tanh \frac{h}{2} = 4\pi \left(\sum_r \delta(z - z_r^+) - \sum_r \delta(z - z_r^-) \right)$$

- Consider $(1, 1)$ vortex pairs on $\Sigma = \mathbb{C}$

Solving the (1,1) Taubes equation (numerically)

$$\nabla^2 h - 2 \tanh \frac{h}{2} = 4\pi (\delta(z - \varepsilon) - \delta(z + \varepsilon))$$

- Regularize: $h = \log \left(\frac{|z-\varepsilon|^2}{|z+\varepsilon|^2} \right) + \hat{h}$


$$\nabla^2 \hat{h} - 2 \frac{|z - \varepsilon|^2 e^{\hat{h}} - |z + \varepsilon|^2}{|z - \varepsilon|^2 e^{\hat{h}} + |z + \varepsilon|^2} = 0$$

- Rescale: $z =: \varepsilon w$

$$\nabla_w^2 \hat{h} - 2\varepsilon^2 \frac{|w - 1|^2 e^{\hat{h}} - |w + 1|^2}{|w - 1|^2 e^{\hat{h}} + |w + 1|^2} = 0$$


- Solve with b.c. $\hat{h}(\infty) = 0$

(1,1) vortices




$$\varepsilon = 4$$

(1,1) vortices




$$\varepsilon = 2$$

(1,1) vortices




$$\varepsilon = 1$$

(1,1) vortices




$$\varepsilon = 0.5$$

(1,1) vortices




$$\varepsilon = 0.3$$

(1,1) vortices



$$\varepsilon = 0.15$$

(1,1) vortices



$$\varepsilon = 0.06$$

The metric on M_{n_+, n_-}

- Restriction of kinetic energy

$$T = \frac{1}{2} \int_{\Sigma} (|\dot{\mathbf{n}}|^2 + |\dot{\mathbf{A}}|^2)$$

to M_{n_+, n_-} equips it with a Riemannian metric

- Expand solution h of Taubes eqn about \pm vortex position z_s :

$$\pm h = \log |z - z_s|^2 + a_s + \frac{1}{2} \bar{b}_s(z - z_s) + \frac{1}{2} b_s(\bar{z} - \bar{z}_s) + \dots$$

- $b_r(z_1, \dots, z_{n_+ + n_-})$ (unknown) complex functions
- **Proposition** (Romão-JMS, following Strachan-Samols):

$$g = 2\pi \left\{ \sum_r \Omega(|z_r|) |dz_r|^2 + \sum_{r,s} \frac{\partial b_s}{\partial z_r} dz_r d\bar{z}_s \right\}$$

Holds on any Riemann surface (including \mathbb{C})

The metric on $M_{1,1}(\mathbb{C})$


- $M_{1,1} = (\mathbb{C} \times \mathbb{C}) \setminus \Delta = \mathbb{C}_{com} \times \mathbb{C}^\times$
- $M_{1,1}^0 = \mathbb{C}^\times$

$$g^0 = 2\pi \left(2 + \frac{1}{\varepsilon} \frac{d}{d\varepsilon}(\varepsilon b(\varepsilon)) \right) (d\varepsilon^2 + \varepsilon^2 d\psi^2)$$

where $b(\varepsilon) = b_+(\varepsilon, -\varepsilon)$


- $\varepsilon b(\varepsilon) = \frac{\partial \widehat{h}}{\partial w_1} \Big|_{w=1} - 1$
- Can easily extract this from our numerics

The metric on $M_{1,1}(\mathbb{C})$




$$\varepsilon b(\varepsilon) = \left. \frac{\partial \hat{h}}{\partial w_1} \right|_{w=1} - 1$$

The metric on $M_{1,1}(\mathbb{C})$



$$F(\varepsilon) = 2\pi \left(2 + \frac{1}{\varepsilon} \frac{d(\varepsilon b(\varepsilon))}{d\varepsilon} \right)$$


The metric on $M_{1,1}(\mathbb{C})$: conjectured asymptotics



$$F_*(\varepsilon) = 2\pi(2 + 4K_0(\varepsilon) - 2\varepsilon K_1(\varepsilon)) \sim -8\pi \log \varepsilon$$


$$F_\infty(\varepsilon) = 2\pi \left(2 + \frac{q^2}{\pi^2} K_0(2\varepsilon) \right)$$

Self similarity as $\varepsilon \rightarrow 0$




- Suggests $\widehat{h}_\varepsilon(w) \approx \varepsilon f_*(\varepsilon w)$ for small ε , where f_* is fixed?
- Define $f_\varepsilon(z) := \varepsilon^{-1} \widehat{h}_\varepsilon(\varepsilon^{-1} z)$


Self similarity as $\varepsilon \rightarrow 0$




Self similarity as $\varepsilon \rightarrow 0$



Self similarity as $\varepsilon \rightarrow 0$



Self similarity as $\varepsilon \rightarrow 0$



Self similarity as $\varepsilon \rightarrow 0$

$$(\nabla^2 \hat{h})(w) = 2\varepsilon^2 \frac{|w - 1|^2 e^{\hat{h}(w)} - |w + 1|^2}{|w - 1|^2 e^{\hat{h}(w)} + |w + 1|^2}$$

Self similarity as $\varepsilon \rightarrow 0$

$$(\nabla^2 \hat{h})(w) = 2\varepsilon^2 \frac{|w - 1|^2 e^{\hat{h}(w)} - |w + 1|^2}{|w - 1|^2 e^{\hat{h}(w)} + |w + 1|^2}$$

- Subst $\hat{h}(w) = \varepsilon f_\varepsilon(\varepsilon w)$

Self similarity as $\varepsilon \rightarrow 0$

$$(\nabla^2 f_\varepsilon)(z) = \frac{2}{\varepsilon} \frac{|z - \varepsilon|^2 e^{\varepsilon f_\varepsilon(z)} - |z + \varepsilon|^2}{|z - \varepsilon|^2 e^{\varepsilon f_\varepsilon(z)} + |z + \varepsilon|^2}$$

- Subst $\hat{h}(w) = \varepsilon f_\varepsilon(\varepsilon w)$

Self similarity as $\varepsilon \rightarrow 0$

$$(\nabla^2 f_\varepsilon)(z) = \frac{2}{\varepsilon} \frac{|z - \varepsilon|^2 e^{\varepsilon f_\varepsilon(z)} - |z + \varepsilon|^2}{|z - \varepsilon|^2 e^{\varepsilon f_\varepsilon(z)} + |z + \varepsilon|^2}$$

- Subst $\hat{h}(w) = \varepsilon f_\varepsilon(\varepsilon w)$
- Take formal limit $\varepsilon \rightarrow 0$

Self similarity as $\varepsilon \rightarrow 0$

$$(\nabla^2 f_*)(z) = f_*(z) - \frac{2(z + \bar{z})}{|z|^2}$$

- Subst $\hat{h}(w) = \varepsilon f_\varepsilon(\varepsilon w)$
- Take formal limit $\varepsilon \rightarrow 0$

Self similarity as $\varepsilon \rightarrow 0$

$$(\nabla^2 f_*)(z) = f_*(z) - \frac{2(z + \bar{z})}{|z|^2}$$

- Subst $\hat{h}(w) = \varepsilon f_\varepsilon(\varepsilon w)$
- Take formal limit $\varepsilon \rightarrow 0$
- Screened inhomogeneous Poisson equation, source $-4 \cos \theta / r$


Self similarity as $\varepsilon \rightarrow 0$

$$(\nabla^2 f_*)(z) = f_*(z) - \frac{2(z + \bar{z})}{|z|^2}$$

- Subst $\hat{h}(w) = \varepsilon f_\varepsilon(\varepsilon w)$
- Take formal limit $\varepsilon \rightarrow 0$
- Screened inhomogeneous Poisson equation, source $-4 \cos \theta / r$
- Unique solution (decaying at infinity)

$$f_*(r e^{i\theta}) = \frac{4}{r} (1 - r K_1(r)) \cos \theta$$

Self similarity as $\varepsilon \rightarrow 0$



The metric on $M_{1,1}^0$


- Predict, for small ε ,

$$\widehat{h}(w_1 + i0) \approx \varepsilon f_*(\varepsilon w_1) = \frac{4}{w_1} (1 - \varepsilon w_1 K_1(\varepsilon w_1))$$


whence we extract predictions for $\varepsilon b(\varepsilon)$, $F(\varepsilon)$

$$g^0 = F(\varepsilon)(d\varepsilon^2 + \varepsilon^2 d\psi^2)$$

- Conjecture: $F(\varepsilon) \sim -8\pi \log \varepsilon$ as $\varepsilon \rightarrow 0$
- $M_{1,1}$ is **incomplete**, with unbounded curvature



Vortices on S^2 : $M_{1,1}(S^2)$



- $M_{1,1} = S^2 \times S^2 \setminus \Delta = (0, 1) \times SO(3) \sqcup \{1\} \times S^2$
- g is $SO(3)$ -invariant, kähler, and invariant under $(z_+, z_-) \mapsto (z_-, z_+)$
- Every such metric takes the form

$$g = -\frac{Q'(\varepsilon)}{\varepsilon}(d\varepsilon^2 + \varepsilon^2 \sigma_3^2) + Q(\varepsilon) \left(\frac{1-\varepsilon^2}{1+\varepsilon^2} \sigma_1^2 + \frac{1+\varepsilon^2}{1-\varepsilon^2} \sigma_2^2 \right),$$

for $Q : (0, 1] \rightarrow [0, \infty)$ decreasing with $Q(1) = 0$.

Vortices on S^2 : $M_{1,1}(S^2)$

- $\text{Vol}(M_{1,1}(S^2))$ is finite iff $Q : (0, 1] \rightarrow [0, \infty)$ is bounded

$$\text{Vol}(M_{1,1}(S^2)) = \left[\lim_{\varepsilon \rightarrow 0} 2\pi Q(\varepsilon) \right]^2$$

- How do we extract $Q(\varepsilon)$? Taubes/localization

$$\nabla^2 h - \frac{8R^2}{(1 + |z|^2)^2} \tanh \frac{h}{2} = 4\pi (\delta(z - \varepsilon) - \delta(z + \varepsilon))$$

where $h = \log((1 - \mathbf{e} \cdot \mathbf{n})/(1 + \mathbf{e} \cdot \mathbf{n}))$

- $\pm h = \log |z - z_s|^2 + a_s + \frac{1}{2}\bar{b}_s(z - z_s) + \frac{1}{2}b_s(\bar{z} - \bar{z}_s) + \dots$

$$g = 2\pi \left\{ \sum_r \frac{4R^2}{(1 + |z_r|^2)^2} |dz_r|^2 + \sum_{r,s} \frac{\partial b_s}{\partial z_r} dz_r d\bar{z}_s \right\}$$

The (1,1) Taubes equation on S^2

$$\nabla^2 h - \frac{8R^2}{(1 + |z|^2)^2} \tanh \frac{h}{2} = 4\pi (\delta(z - \varepsilon) - \delta(z + \varepsilon))$$

The (1,1) Taubes equation on S^2

$$\nabla^2 h - \frac{8R^2}{(1+|z|^2)^2} \tanh \frac{h}{2} = 4\pi (\delta(z-\varepsilon) - \delta(z+\varepsilon))$$

- Regularize: $h = \log \left(\frac{|z-\varepsilon|^2}{|z+\varepsilon|^2} \right) + \hat{h}$

The (1,1) Taubes equation on S^2

$$\nabla^2 h - \frac{8R^2}{(1+|z|^2)^2} \tanh \frac{h}{2} = 4\pi (\delta(z-\varepsilon) - \delta(z+\varepsilon))$$

- Regularize: $h = \log \left(\frac{|z-\varepsilon|^2}{|z+\varepsilon|^2} \right) + \hat{h}$
- Rescale: $z =: \varepsilon w$

The (1,1) Taubes equation on S^2

$$\nabla_w^2 \hat{h} - \frac{8R^2\varepsilon^2}{(1+\varepsilon^2|w|^2)^2} \frac{|w-1|^2 e^{\hat{h}} - |w+1|^2}{|w-1|^2 e^{\hat{h}} + |w+1|^2} = 0$$

- Regularize: $h = \log\left(\frac{|z-\varepsilon|^2}{|z+\varepsilon|^2}\right) + \hat{h}$
- Rescale: $z =: \varepsilon w$

The (1,1) Taubes equation on S^2

$$-\Delta_{S^2} \hat{h} + 8R^2 \varepsilon^2 \left(\frac{1 + |w|^2}{1 + \varepsilon^2 |w|^2} \right)^2 \frac{|w - 1|^2 e^{\hat{h}} - |w + 1|^2}{|w - 1|^2 e^{\hat{h}} + |w + 1|^2} = 0$$

- Regularize: $h = \log \left(\frac{|z - \varepsilon|^2}{|z + \varepsilon|^2} \right) + \hat{h}$
- Rescale: $z =: \varepsilon w$

The (1,1) Taubes equation on S^2

$$-\Delta_{S^2} \hat{h} + 8R^2 \varepsilon^2 G(\varepsilon, w) F(w, \hat{h}) = 0$$

- Regularize: $h = \log\left(\frac{|z-\varepsilon|^2}{|z+\varepsilon|^2}\right) + \hat{h}$
- Rescale: $z =: \varepsilon w$
- $G : \mathbb{R} \times S^2 \rightarrow \mathbb{R}$, $F : S^2 \times \mathbb{R} \rightarrow \mathbb{R}$ smooth

The (1,1) Taubes equation on S^2

$$-\Delta_{S^2} \hat{h} + 8R^2 \varepsilon^2 G(\varepsilon, w) F(w, \hat{h}) = 0$$

- Regularize: $h = \log\left(\frac{|z-\varepsilon|^2}{|z+\varepsilon|^2}\right) + \hat{h}$
- Rescale: $z =: \varepsilon w$
- $G : \mathbb{R} \times S^2 \rightarrow \mathbb{R}$, $F : S^2 \times \mathbb{R} \rightarrow \mathbb{R}$ smooth
- $\varepsilon b(\varepsilon) = \hat{h}_x(1, 0) - 1$

The (1,1) Taubes equation on S^2

$$-\Delta_{S^2} \hat{h} + 8R^2 \varepsilon^2 G(\varepsilon, w) F(w, \hat{h}) = 0$$

- Regularize: $h = \log\left(\frac{|z-\varepsilon|^2}{|z+\varepsilon|^2}\right) + \hat{h}$
- Rescale: $z =: \varepsilon w$
- $G : \mathbb{R} \times S^2 \rightarrow \mathbb{R}$, $F : S^2 \times \mathbb{R} \rightarrow \mathbb{R}$ smooth
- $\varepsilon b(\varepsilon) = \hat{h}_x(1, 0) - 1$
- $Q(\varepsilon) = -2\pi \left(1 + 2R^2 + \varepsilon b(\varepsilon) - \frac{4R^2}{1 + \varepsilon^2}\right)$

The regularized (1,1) Taubes equation on S^2

$$-\Delta_{S^2} \hat{h} + 8R^2 \varepsilon^2 G(\varepsilon, w) F(w, \hat{h}) = 0 \quad (*)$$

- Elliptic estimates
- $\tilde{h}(z) = \hat{h}(z/\varepsilon)$, $*F_A = -\frac{1}{2}\Delta_{S^2} \tilde{h}$

$$\|\tilde{h}\|_{H^2(S^2)}^2 \leq C \|\Delta_{S^2} \tilde{h}\|_{L^2}^2 = C \|F_A\|_{L^2}^2 \leq C$$

- Sobolev: $\|\hat{h}\|_{C^0} = \|\tilde{h}\|_{C^0} \leq C \|\tilde{h}\|_{H^2} \leq C$
- Allows us to prove more refined estimate

$$\|\hat{h}\|_{H^1(S^2)}^2 \leq C \langle \hat{h}, -\Delta_{S^2} \hat{h} \rangle \leq C\varepsilon$$

whence (diff $(*)$ wrt x , estimate on a disk around $(1, 0)$)

$$\|\partial_x \hat{h}\|_{H^2(\mathbb{D})} \leq C\varepsilon$$

- Sobolev again: $\|\hat{h}_x\|_{C^0(\mathbb{D})} \leq C\varepsilon^{1/2} \Rightarrow |\partial_x \hat{h}(1, 0)| \leq C\varepsilon^{1/2}$

The regularized (1,1) Taubes equation on S^2

- Hence $\lim_{\varepsilon \rightarrow 0} \varepsilon b(\varepsilon) = -1$
- Hence $\lim_{\varepsilon \rightarrow 0} Q(\varepsilon) = 4\pi R^2$
- **Theorem** (Romão, JMS) Let Σ be a round two-sphere. Then

$$Vol(M_{1,1}(\Sigma)) = (2\pi Vol(\Sigma))^2.$$

Completeness?

- Need to know length of radial geodesic $0 < \varepsilon \leq 1$ in $M_{1,1}(S^2)$
- $M_{1,1}^0(S^2) = S^2 \setminus \{(0, 0, 1)\}$

$$g^0 = F(\varepsilon)(d\varepsilon^2 + \varepsilon^2 d\psi^2)$$

- $L = \int_0^1 \sqrt{F(\varepsilon)} d\varepsilon$
- $F(\varepsilon)/2\pi = \frac{8R^2}{(1+\varepsilon^2)^2} + \frac{1}{\varepsilon} \frac{d(\varepsilon b(\varepsilon))}{d\varepsilon}$
- $\varepsilon b(\varepsilon) + 1 = \partial_x \hat{h}(0, 1)$
- Need to control $\partial_x \partial_\varepsilon \hat{h}$ at $(0, 1)$

The regularized (1,1) Taubes equation on S^2

$$-\Delta_{S^2} \hat{h}_\varepsilon + 8R^2 \varepsilon^2 G(\varepsilon, w) F(w, \hat{h}_\varepsilon) = 0 \quad (*)$$

- $u_\varepsilon := \partial \hat{h}_\varepsilon / \partial \varepsilon : S^2 \rightarrow \mathbb{R}$

$$-\Delta_{S^2} u_\varepsilon + G_1(\varepsilon, w, \hat{h}_\varepsilon) u_\varepsilon + G_2(\varepsilon, w, \hat{h}_\varepsilon) = 0$$

- Elliptic estimates: $\|\partial_x u_\varepsilon\|_{H^2(\mathbb{D})} \leq C\varepsilon^{-1/2}$
- Conformal factor

$$\frac{F(\varepsilon)}{2\pi} = \frac{8R^2}{(1+\varepsilon^2)^2} + \frac{1}{\varepsilon} \partial_x u_\varepsilon \Big|_{(1,0)} \leq C(1 + \varepsilon^{-3/2})$$

- Radial geodesic has length $L = \int_0^1 \sqrt{F(\varepsilon)} d\varepsilon < \infty$
- **Theorem (Romão, JMS)** $M_{1,1}(S^2)$ is geodesically incomplete

The volume of $M_{n,n}(S^2)$

- $M_{n,n}(S^2) = \{\text{disjoint pairs of } n\text{-divisors on } S^2\} = (\mathbb{P}^n \times \mathbb{P}^n) \setminus \Delta$
- Consider gauged **linear** sigma model:
 - fibre \mathbb{C}^2
 - gauge group $\tilde{U}(1) \times U(1) : (\varphi_1, \varphi_2) \mapsto (e^{i(\tilde{\theta}+\theta)} \varphi_1, e^{i\tilde{\theta}} \varphi_2)$

$$\begin{aligned} E_{\tilde{e}} &= \frac{1}{2} \int_{\Sigma} \left\{ \frac{|\tilde{F}|^2}{\tilde{e}^2} + |F|^2 + |\mathrm{d}_{\tilde{A}}\varphi|^2 + |\mathrm{d}_A\varphi|^2 \right. \\ &\quad \left. + \frac{\tilde{e}^2}{4} (4 - |\varphi_1|^2 - |\varphi_2|^2)^2 + \frac{1}{4} (2 - |\varphi_1|^2)^2 \right\} \end{aligned}$$

- For any $\tilde{e} > 0$, has compact moduli space of (n, n) -vortices

$$M_{n,n}^{\text{lin}} = \mathbb{P}^n \times \mathbb{P}^n$$

- Baptista found a formula for $[\omega_{L^2}]$ of $M_{n_1, n_2}^{\text{lin}}(\Sigma)$
- Can compute $\text{Vol}(M_{n,n}^{\text{lin}}(S^2))$ by evaluating $[\omega_{L^2}]$ on $\mathbb{P}^1 \times \{p\}$, $\{p\} \times \mathbb{P}^1$

The volume of $M_{n,n}(S^2)$

$$\begin{aligned} E_{\tilde{e}} = & \frac{1}{2} \int_{\Sigma} \left\{ \frac{|\tilde{F}|^2}{\tilde{e}^2} + |F|^2 + |\mathrm{d}_{\tilde{A}}\varphi|^2 + |\mathrm{d}_A\varphi|^2 \right. \\ & \left. + \frac{\tilde{e}^2}{4} (4 - |\varphi_1|^2 - |\varphi_2|^2)^2 + \frac{1}{4} (2 - |\varphi_1|^2)^2 \right\} \end{aligned}$$

- Take formal limit $\tilde{e} \rightarrow 0$:
 - $|\varphi_1|^2 + |\varphi_2|^2 = 4$ pointwise
 - \tilde{A} frozen out, fibre \mathbb{C}^2 collapses to $S^3/\widetilde{U}(1) = \mathbb{P}^1$
 - E-L eqn for \tilde{A} is algebraic: eliminate \tilde{A} from E_∞

$$E_\infty = \frac{1}{2} \int_{\Sigma} |F|^2 + 4 \frac{|\mathrm{d}u - iAu|^2}{(1 + |u|^2)^2} + \left(\frac{1 - |u|^2}{1 + |u|^2} \right)^2$$

where $u = \varphi_1/\varphi_2$

- Exactly our \mathbb{P}^1 sigma model!

The volume of $M_{n,n}(S^2)$

- Leads us to conjecture that

$$\text{Vol}(M_{n,n}(S^2)) = \lim_{\tilde{\epsilon} \rightarrow \infty} \text{Vol}(M_{n,n}^{\text{lin}}(S^2)) = \frac{(2\pi \text{Vol}(S^2))^{2n}}{(n!)^2}$$

Agrees with $M_{1,1}(S_R^2)$.

- Can generalize to $n_+ > n_-$:

$$\text{Vol}(M_{n,m}(S^2)) = \frac{(2\pi)^{n+m}}{n!m!} (\text{Vol}(S^2) - \pi(n-m))^n (\text{Vol}(S^2) + \pi(n-m))^m$$

and to $\text{genus}(\Sigma) > 0$

- Similar conjectures for Einstein-Hilbert action...
- Similar limit (\mathbb{C}^k fibre, $U(1)$ gauge \rightarrow ungauged \mathbb{P}^{k-1} model) studied rigorously by Chih-Chung Liu.
- Thermodynamics of vortex gas **mixture**

Summary / What next?

- Case $\Sigma = \mathbb{C}$ is most interesting
- $M_{1,1}(\mathbb{C}) = \mathbb{C} \times \mathbb{C} \setminus \Delta = \mathbb{C}_{\text{com}} \times \mathbb{C}^\times$
- Numerics: metric on SoR \mathbb{C}^\times , $g^0 = F(\varepsilon)(d\varepsilon^2 + \varepsilon^2 d\psi^2)$
- Conjectured asymptotics in small ε region

$$F(\varepsilon) \sim -8\pi \log \varepsilon$$

- Would imply $M_{1,1}(\mathbb{C})$ is incomplete with unbounded scalar curvature
- Can we prove it?
- We can shift the vacuum manifold:

$$\mu(\mathbf{n}) = \tau - \mathbf{e} \cdot \mathbf{n}$$

Case $0 < \tau < 1$ very sparsely explored